arXiv:1212.2712v1 [physics.flu-dyn] 12 Dec 2012

A General Method for Large Eddy Simulation in Different Range of Turbulent Flows

Changping Yu[1]

[1] *Peking University, Beijing 100871, China*

(Dated: December 13, 2012)

Abstract

Traditional large eddy simulation is based on Kolmogrov's hypothesis, and done in the inertial range. In inertial range the LES model coefficient is scale-invariant. In many cases, such as computing in the boundary layer, the filter scale is not in the inertial range, and the model coefficient changes drastically with the filter scale, thus the assumption of scale-invariance is not proper. In this paper, we propose a general principle of constraint, and then we deduce two concrete constraints based on the existing spectrum theory and priori data. The constrained dynamic Smogorinsky model can predict energy spectrum well in different range of forced and decayed turbulence. Dynamic mixed models with one of the constraints shows more efficient than some classical models in predicting energy spectrum and SGS dissipation, and also displays a strong correlation with the real stress in different range of the turbulent flows.

PACS numbers:

I. INTRODUCTION

Large eddy simulation (LES) is an important method in numerical simulation of turbulent flows [1–5]. Traditional large eddy simulation is based on the K41 theory (Kolmogrov,1941), and done in the inertial range. In the inertial range, the turbulent motions are statistically isotropic, and the statistics of the motions are decided by the dissipation rate and wave number .When doing LES, the motions larger than filter scale are solved directly, and the motions smaller than filter scale (SGS) are modeled by the resolved quantities. Several kinds of SGS models have been proposed so far, such as eddy-viscosity model, dynamic model, vortex model, etc.[6, 7] Recently, a type of constrained LES method has already been developed [8]. The traditional SGS model fits the assumption of scale-invariance in the inertial range. Dynamic Smogorinsky model (DSM)is a classic model in large eddy simulation of turbulent flows [5]. Here dynamic procedure is based on the Germano identity,

$$L_{ij} = T_{ij} - \overline{\tau}_{ij} = \overline{\widetilde{u}_i \widetilde{u}_j} - \overline{\widetilde{u}_i}\ \overline{\widetilde{u}_j}, \tag{1}$$

Where L_{ij} is called the resolved stress, $\tau_{ij} = \widetilde{u_i u_j} - \widetilde{u}_i \widetilde{u}_j$ is the SGS stress at the filtered scale Δ, and $T_{ij} = \overline{\widetilde{u_i u_j}} - \overline{\widetilde{u}}_i \overline{\widetilde{u}}_j$ is the SGS stress at the filtered scale $\alpha\Delta(\alpha > 1)$. Here a tilde denotes filtering at scale Δ and a bar denotes test-filtering at scale at $\alpha\Delta$. In (1), taking Smogorinsky model stress τ_{ij}^{mod} and T_{ij}^{mod} to replace τ_{ij} and T_{ij}, we can get

$$L_{ij}^{\mathrm{mod}} = T_{ij}^{\mathrm{mod}} - \overline{\tau_{ij}^{\mathrm{mod}}}, \tag{2}$$

Where L_{ij}^{mod} is called the resolved stress,

$$\tau_{ij}^{\mathrm{mod}} = C_{sm}\Delta^2 |\widetilde{S}| \widetilde{S}_{ij}, \tag{3}$$

$$T_{ij}^{\mathrm{mod}} = C_{sm}(\alpha\Delta)^2 |\overline{\widetilde{S}}| \overline{\widetilde{S}}_{ij}, \tag{4}$$

and here $\widetilde{S}_{ij}$ is resolved strain tensor at filtering scale Δ, $|\widetilde{S}| = \left(2\widetilde{S}_{ij}\widetilde{S}_{ij}\right)^{1/2}$. A square error between L_{ij} and L_{ij}^{mod} can be given by

$$E^{\mathrm{mod}} = \langle (L_{ij} - L_{ij}^{\mathrm{mod}})^2 \rangle, \tag{5}$$

Here the bracket $\langle \cdot \rangle$ denotes ensemble average of the turbulence flow. According to the hypothesis of scale-invariance in the inertial range of turbulence flow, the model coefficients are same in τ_{ij}^{mod} and T_{ij}^{mod}. Minimizing (5), we can get the expression of the model coefficients,

$$C_{sm} = \frac{\langle M_{ij} L_{ij} \rangle}{\langle M_{ij} M_{ij} \rangle}, \tag{6}$$

where,

$$M_{ij} = (\alpha\Delta)^2|\overline{\widetilde{S}}|\overline{\widetilde{S}}_{ij} - \Delta^2\overline{|\widetilde{S}|\widetilde{S}_{ij}}, \tag{7}$$

When do LES using the dynamic Smogorinsky model, the test-scale and even the filter-scale are often out of the inertial range. Specially, more and more researches pay attention to the behavior of turbulence in dissipation range [9–11]. Thus the assumption of scale-invariance is not proper yet [12], and the validity of the traditional LES result is also not credible. In order to solve this problem, several methods have been proposed so far. Ronchi et al. have proposed a complex dynamic model [13], and the model coefficient varies with scale. Although the practicability of the model has not been tested widely, it supplies a feasible idea to us all. Using several existing energy spectrum, P. R. Voke gives the approximative coefficient of Smogorinsky model in dissipation range [13]. Recently, Meneveau et al. have proposed a scale-dependent dynamic Smogorinsky model (SDSM) for large-eddy simulation in a series of articles [14–19]. They construct a function of Smogorinsky model coefficients at the filter and test-filter scales,

$$f(\Delta) = C_{sm}(\alpha\Delta)/C_{sm}(\Delta), \tag{8}$$

where the expression of $f\left(\Delta\right)$ can be get through priori knowledge. Thus (7) can be rewrite as

$$M_{ij} = f(\Delta)(\alpha\Delta)^2|\overline{\widetilde{S}}|\overline{\widetilde{S}}_{ij} - \Delta^2\overline{|\widetilde{S}|\widetilde{S}_{ij}}, \tag{9}$$

Taking (9) into (6) , we can get the Smogorinsky model coefficient in the dissipation range.

In this paper, we first propose a general constraint principle to construct the relation between two different scales. Then we give two concrete constraints through energy spectrum and priori knowledge. It has been developed several types of energy spectrum so far [20–23]. In this paper we choose Kovasznay spectrum [24], and apply the two constrained condition to single model and mixed models [26–28]. The results of LES with any one of the two constraints show some superiority than other models.

II. THE GENERAL CONSTRAINT

As is well known that traditional LES is done in the inertial range, and the SGS dissipation ε^{SGS} equals to total dissipation approximately. In general, total dissipation is made up of two parts, the SGS dissipation and the viscous dissipation ε^{ν} at the filter scale of LES,

$$\varepsilon = \varepsilon^{\text{SGS}} + \varepsilon^{\nu}. \tag{10}$$

And we have an identical equation on SGS dissipation at scale Δ and $\alpha\Delta$,

$$\frac{\langle \tau_{ij}\widetilde{S}_{ij}\rangle}{\varepsilon_{\Delta}^{\mathrm{SGS}}} = \frac{\langle T_{ij}\overline{\widetilde{S}}_{ij}\rangle}{\varepsilon_{(}\alpha\Delta)^{\mathrm{SGS}}}. \tag{11}$$

Then we can construct the relation of $\varepsilon^{\mathrm{SGS}}$ and ε^{ν},

$$\varepsilon^{\mathrm{SGS}} = F(\varepsilon^{\nu}). \tag{12}$$

Thus, putting (12) into (11), we can get the identical equation as following,

$$\frac{\langle \tau_{ij}\widetilde{S}_{ij}\rangle}{F(\varepsilon_{\Delta}^{\nu})} = \frac{\langle T_{ij}\overline{\widetilde{S}}_{ij}\rangle}{F(\varepsilon_{\alpha\Delta}^{\nu})}, \tag{13}$$

Taking model stress τ_{ij}^{mod} and T_{ij}^{mod} to replace τ_{ij} and T_{ij}, we can get

$$\frac{\langle \tau_{ij}^{\mathrm{mod}}\widetilde{S}_{ij}\rangle}{F(\varepsilon_{\Delta}^{\nu})} = \frac{\langle T_{ij}^{\mathrm{mod}}\overline{\widetilde{S}}_{ij}\rangle}{F(\varepsilon_{\alpha\Delta}^{\nu})}. \tag{14}$$

Equation (14) is the general principle of constraint. Next, we will give two concrete form of $F\left(\varepsilon^{\nu}\right)$ based on the existing spectrum theory and priori data.

A. Constrained condition based on spectrum theory

Kovasznay (1948) proposed a energy spectrum, and it supports a energy distribution with wave number from inertial range to dissipation range. The form of Kovasznay spectrum is

$$E(k) = C_k\varepsilon^{2/3}k^{-5/3}\left(1-\frac{C_k}{2}\left(\frac{k}{k_d}\right)^{4/3}\right), \quad k_d = (\frac{\varepsilon}{\nu^3})^{1/4}. \tag{15}$$

The viscous dissipation at filter scale Δ can be expressed as

$$\varepsilon_{\Delta}^{\nu} = 2\nu\int_0^{k_c} k^2E(k)\,dk, \quad k_c = \pi/\Delta. \tag{16}$$

From (15) and (16) we know that

$$\varepsilon_{\Delta}^{\nu} = \nu k_d^{4/3}\varepsilon^{2/3}\left(1-\left(1-\frac{C_k}{2}\left(\frac{k_c}{k_d}\right)^{4/3}\right)^3\right). \tag{17}$$

At filter scale Δ, (10) can be written as

$$\varepsilon = \varepsilon_{\Delta}^{\mathrm{SGS}} + \varepsilon_{\Delta}^{\nu}. \tag{18}$$

Thus,

$$\frac{\varepsilon_{\Delta}^{\nu}}{\varepsilon} = \frac{\varepsilon_{\Delta}^{\nu}}{\varepsilon_{\Delta}^{\nu}+\varepsilon_{\Delta}^{\mathrm{SGS}}} = \frac{1}{1+\eta_{\Delta}}, \quad \eta_{\Delta} = \frac{\varepsilon_{\Delta}^{\mathrm{SGS}}}{\varepsilon_{\Delta}^{\nu}}. \tag{19}$$

From (15), (17) and (19), we can get

$$\frac{\varepsilon_{\Delta}^{\mathrm{SGS}}}{\varepsilon_{\Delta}^{\nu}} = 1 - \left(1 - \frac{C_k}{2}\left(\frac{k_c}{k_d}\right)^{4/3}\right)^3, \tag{20}$$

where,

$$\begin{aligned}
\frac{k_c}{k_d} = \frac{\pi}{\Delta}\left(\frac{\nu^3}{\varepsilon}\right)^{1/4} &= \frac{\pi}{\Delta}\left(\frac{\nu^2 \cdot 2\nu\langle\widetilde{S}_{ij}\widetilde{S}_{ij}\rangle}{2\langle\widetilde{S}_{ij}\widetilde{S}_{ij}\rangle}\right)^{1/4} \\
&= \frac{\pi\nu^{1/2}}{\Delta\left(2\langle\widetilde{S}_{ij}\widetilde{S}_{ij}\rangle\right)^{1/4}}\left(\frac{\varepsilon_{\Delta}^{\nu}}{\varepsilon}\right)^{1/4} \\
&= \frac{\pi\nu^{1/2}}{\Delta\left(2\langle\widetilde{S}_{ij}\widetilde{S}_{ij}\rangle\right)^{1/4}}\left(1+\eta_{\Delta}\right)^{-1/4}. \qquad (21)
\end{aligned}$$

Putting (21) into (20), the result can be got

$$\begin{aligned}
\frac{1}{1+\eta_{\Delta}} &= 1 - \left[1 - \beta_{\Delta}\left(1+\eta_{\Delta}\right)^{-1/3}\right]^3, \\
\beta_{\Delta} &= \frac{C_k}{2}\frac{\pi^{4/3}\nu^{2/3}}{\Delta^{4/3}\left(2\langle\widetilde{S}_{ij}\widetilde{S}_{ij}\rangle\right)^{1/3}}. \qquad (22)
\end{aligned}$$

From (22), we know

$$\eta_{\Delta} = \frac{1}{8}\left(\beta_{\Delta} + \sqrt{\frac{4-3\beta_{\Delta}^3}{3\beta_{\Delta}}}\right)^3 - 1. \tag{23}$$

Therefore, the expression of $F\left(\varepsilon^{\nu}\right)$ can be written as

$$F\left(\varepsilon_{\Delta}^{\nu}\right) = \eta_{\Delta}\varepsilon_{\Delta}^{\nu}. \tag{24}$$

In general case, the equation can be got approximately that

$$\overline{S}_{ij} = \overline{\widetilde{S}}_{ij}, \tag{25}$$

where, $\widetilde{}$ denotes filtering at Δ and at scale $\alpha\Delta$. Thus, the a concrete form (14) can be expressed as

$$\frac{\langle\tau_{ij}^{\mathrm{mod}}\widetilde{S}_{ij}\rangle}{\eta_{\Delta}\varepsilon_{\Delta}^{\nu}} = \frac{\langle T_{ij}^{\mathrm{mod}}\overline{\widetilde{S}}_{ij}\rangle}{\eta_{\alpha\Delta}\varepsilon_{\alpha\Delta}^{\nu}}. \tag{26}$$

B. Constrained condition based on priori data fitting

As described by Frisch et al. (1978), the β model quantifies the course of cascade [21]. Similar with this theory, we can propose a supposition on energy dissipation of cascade. On some scale, the viscous dissipation ε^{ν} denotes the loss of kinetic energy and the SGS dissipation $\varepsilon^{\mathrm{SGS}}$ denotes the kinetic energy transferring into the next scale. Based on the scaling similarity of eddy structure, $\varepsilon^{\nu}_{\Delta}/\varepsilon^{\mathrm{SGS}}_{\Delta}$ can be expressed by the function of Δ (Δ is non-dimensional number of filter scale)

$$\varepsilon^{\nu}_{\Delta}/\varepsilon^{\mathrm{SGS}}_{\Delta} = f(\Delta). \tag{27}$$

As $\Delta \to 0$, we know that $\varepsilon^{\mathrm{SGS}}_{\Delta} \to 0$ and $\varepsilon^{\nu}_{\Delta} \to c$ (constant). While Δ is in the inertial range even the energy-containing range, $\varepsilon^{\nu}_{\Delta}$ is ignorable in comparison with $\varepsilon^{\mathrm{SGS}}_{\Delta}$. Therefore, we can give the expression of $f(\Delta)$, and two-term polynomial is used here,

$$f(\Delta) = C_{\nu}(\Delta^{\gamma 1} + \Delta^{\gamma 2}), \tag{28}$$

where, C_{ν} is the coefficient, and C_{ν} is variable with different kinetic viscous ν. In order to check the hypothesis (28), we can give the result of priori as Fig. 1 From Fig. 1, we can see that $C_{\nu}\varepsilon^{\nu}_{\Delta}/\varepsilon^{\mathrm{SGS}}_{\Delta}$ can be expressed by an unified function. Thus, t proves that the hypothesis (28) is reasonable. Then we choose one of the viscous ν and fit the priori result of $\varepsilon^{\nu}_{\Delta}/\varepsilon^{\mathrm{SGS}}_{\Delta}$. Thus we find the concrete form of γ



FIG. 1: (Color online) In the case of different ν, the distribution of $C_{\nu}\varepsilon^{\nu}_{\Delta}/\varepsilon^{\mathrm{SGS}}_{\Delta}$ with changing scale for priori Line with circle: $\nu = 0.0006$; Line with square: $\nu = 0.001$; Line with delta: $\nu = 0.002$; Line with RTringle: $\nu = 0.005$.

$$\gamma 1 = -2, \gamma 1 = -1. \tag{29}$$

The constrained condition can also be expressed as below

$$\frac{\langle \tau^{\mathrm{mod}}_{ij} \widetilde{S}_{ij} \rangle}{\varepsilon^{\nu}_{\Delta}/(\Delta^{-2} + \Delta^{-1})} = \frac{\langle T^{\mathrm{mod}}_{ij} \overline{\widetilde{S}}_{ij} \rangle}{\varepsilon^{\nu}_{\alpha\Delta}/((\alpha\Delta)^{-2} + (\alpha\Delta)^{-1})}, \tag{30}$$

II A and II B introduce two concrete forms of (14), and then we will introduce a new mix dynamic model used in this paper.

III. THE CONSTRAINED LES MODEL

In this article, we first construct two dynamic Smogorinsky models with constrained condition (26) or (30) (CDSM1, CDSM2) contrast with some single model as dynamic Smogorinsky model and SDSM.

For model CDSM1, we can know that

$$\tau_{ij}^{\mathrm{mod}} = C_{\mathrm{mod}}\Delta^2|\widetilde{S}|\widetilde{S}_{ij}, \tag{31}$$

$$T_{ij}^{\mathrm{mod}} = C_{\mathrm{test}}(\alpha\Delta)^2|\overline{\widetilde{S}}_{ij}|\overline{\widetilde{S}}_{ij}. \tag{32}$$

And the constrained condition is

$$\frac{\langle C_{\mathrm{mod}}\Delta^2|\widetilde{S}|\widetilde{S}_{ij}\widetilde{S}_{ij}\rangle}{\eta_{\Delta}\varepsilon_{\Delta}^{\nu}} = \frac{\langle C_{\mathrm{test}}(\alpha\Delta)^2|\overline{\widetilde{S}}|\overline{\widetilde{S}}_{ij}\overline{\widetilde{S}}_{ij}\rangle}{\eta_{\alpha\Delta}\varepsilon_{\alpha\Delta}^{\nu}}. \tag{33}$$

Thus, we can get the expression of C_{mod},

$$C_{\mathrm{mod}} = \frac{\langle M_{ij}L_{ij}\rangle}{\langle M_{ij}M_{ij}\rangle}, \tag{34}$$

where,

$$M_{ij} = \frac{\langle|\widetilde{S}|\widetilde{S}_{ij}\widetilde{S}_{ij}\rangle\eta_{\alpha\Delta}\varepsilon_{\alpha\Delta}^{\nu}}{\langle|\overline{\widetilde{S}}|\overline{\widetilde{S}}_{ij}\overline{\widetilde{S}}_{ij}\rangle\eta_{\Delta}\varepsilon_{\Delta}^{\nu}}\Delta^2|\overline{\widetilde{S}}|\overline{\widetilde{S}}_{ij} - \Delta^2|\widetilde{S}|\widetilde{S}_{ij}, \tag{35}$$

For model CDSM2,

$$\begin{aligned} M_{ij} \; = \; & \frac{\langle|\widetilde{S}|\widetilde{S}_{ij}\widetilde{S}_{ij}\rangle(\Delta^{-2}+\Delta^{-1})\varepsilon_{\alpha\Delta}^{\nu}}{\langle|\overline{\widetilde{S}}|\overline{\widetilde{S}}_{ij}\overline{\widetilde{S}}_{ij}\rangle(\alpha\Delta)^{-2}+(\alpha\Delta)^{-1})\varepsilon_{\Delta}^{\nu}}\Delta^2|\overline{\widetilde{S}}|\overline{\widetilde{S}}_{ij} \\ & -\Delta^2|\widetilde{S}|\widetilde{S}_{ij}, \end{aligned} \tag{36}$$

Different from other previous methods, the scale-dependent constrained condition can also apply to mix models. Here we propose a constrained dynamic mixed helical SGS models. There are several SGS stress model commonly used in large eddy simulation of turbulence. The Smogorinsky model is a kind of eddy-viscosity model, and it can predict energy dissipation well, and has good numerical robustness and stability. But it can not predict the energy backscatter, and the correlation with the real SGS stress is not good . The nonlinear model, also called gradient or tensor eddy viscosity model, can successfully predict the structure of real stress and energy backscatter and does not need second filtering. Have thought of the advantages and disadvantages of the two models, we can construct a two-term mixed model $\tau_{ij}^{\mathrm{mod}} \cdot \tau_{ij}^{\mathrm{mod}}$ and T_{ij}^{mod} can be expressed below,

$$\tau_{ij}^{\mathrm{mod}} = c_{11}f_{1,ij} + c_{12}f_{2,ij}, \tag{37}$$

$$T_{ij}^{\mathrm{mod}} = c_{21}F_{1,ij} + c_{22}F_{2,ij}, \tag{38}$$

where $f_{1,ij} = \Delta^2|\widetilde{S}|\widetilde{S}_{ij}$, $f_{2,ij} = \Delta^2 \frac{\partial \widetilde{u}_i}{\partial x_k}\frac{\partial \widetilde{u}_j}{\partial x_k}$; $F_{1,ij} = (\alpha\Delta)^2\,|\overline{\widetilde{S}}|\overline{\widetilde{S}}_{ij}$, $F_{2,ij} = (\alpha\Delta)^2\,\frac{\partial \overline{\widetilde{u}}_i}{\partial x_k}\frac{\partial \overline{\widetilde{u}}_j}{\partial x_k}$. Then, with the constrained condition (26) or (30), using (31), we can construct the constrained dynamic mixed models CDMM1 and CDMM2, respectively.

IV. THE ANALYSIS OF THE RESULTS

Now we will give the analysis of the results of the forced steady and decaying homogeneous isotropic turbulence. In order to validate our model, a DNS of three-dimensional incompressible homogeneous isotropic turbulence is introduced here. It solves the forced N-S equations using a pseudo spectral code in a cubic box with periodic boundary conditions, and the numerical resolution is 512^3. A Guassian random field is the initial flow condition, and it has an energy spectrum as,

$$E_0(k) = Ak^2U_0^2k_0^{-5}e^{-\frac{2k^2}{k_0^2}}, \tag{39}$$

where $k_0 = 4.5786$ and $U_0 = 0.715$. The whole system is maintained by a constant energy input rate $\varepsilon = 0.1$ in the first two wave number shells [29].

To validate the constrained condition, we take four types of Smogorinsky models (CDSM1,CDSM2,SDSM and DSM) for comparison firstly. In this paper the LES and DNS of forced turbulence are all performed starting from a Gaussian random field with an inertial energy spectrum as (33), and the kinematic viscosity is $\nu = 0.001$. In Fig. 2, we present the energy spectra of the four type single models for forced steady isotropic turbulence at different filter scales, (a) filter scale $\Delta = 3\pi/64$ (b) filter scale $\Delta = 3\pi/128$. The filter-scale of LES in Fig. 2(a) is close to the inertial range and (b) close to the dissipation range. We can see that the energy spectra of CDSM2 is closer to the DNS spectra than other spectra at most of the wave-number in (a). The energy spectra of CDSM1 and CDSM2 shows better than SDSM and DSM particular in high wave-number, and SDSM is a little better than DSM in (b).

Fig. 3 shows us the evolution of model coefficients (CDSM1, CDSM2, SDSM and DSM) with time in steady turbulence, (a) $\Delta = 3\pi/64$ (b) $\Delta = 3\pi/128$. In Fig. 3, we can see that the distribution of the four model coefficients varied drastically with different filter scale. Near the inertial range (a), the model coefficients of CDSM1, SDSM and DSM are close, and near the dissipation range(b), the model coefficients of CDSM1 and CDSM2 are much closer. Totally, the value of DSM coefficient is smallest, and CDSM2 is biggest.



FIG. 2: (Color online) Energy spectra and for steady forced isotropic turbulence (a posteriori). Solid line: DNS; dashed line: CDSM1; DashDD line: CDSM2; line with diamond: SDSM; line with delta: DSM. (a) filter scale $\Delta = 3\pi/64$ (b) filter scale $\Delta = 3\pi/128$.



FIG. 3: (Color online) model coefficient evolution with time for steady forced isotropic turbulence for a posteriori, (a) at filter-scale $\Delta = 3\pi/64$, (b) at filter-scale $\Delta = 3\pi/128$. Line with square: CDSM1; Line with delta: CDSM2; Line with diamond: SDSM; line with circle: DSM.

Shown in Fig. 4 are the dynamically calculated model and test model coefficients of CDSM1, (a) $\Delta = 3\pi/64$ (b) $\Delta = 3\pi/128$. In Fig. 4, we can see that C_{mod} and C_{test} vary no more than 10% in (a), and while vary about 50%. From these results, we can see that taking C_{mod} and C_{test} the same value is not proper out of inertial range.

Fig. 5 and Fig. 6 show us decay energy spectra of the four model evolution with time ($t = 0$, $6\tau_0$ and $12\tau_0$), starting from a fully developed statistical steady state, (a) $\Delta = 3\pi/64$ (b) $\Delta = 3\pi/128$. Here τ_0 is the initial large eddy turnover time scale, and is defined as $\tau_0 = \pi/(k_0U_0)$. The solid lines are DNS spectra, and dashed lines are energy spectra for the four models. From the two figures, we can see that CDSM1 and CDSM2 display the similar trends and predict the energy spectra very well. And DSM model displays a worse status than other models.

Through the analysis above all, we can say that single model with our constraints (CDSM1 and



FIG. 4: (Color online) model and test coefficients of CDSM1 distribution with time for steady forced isotropic turbulence for a posteriori, (a) at filter-scale $\Delta = 3\pi/64$, (b) at filter-scale $\Delta = 3\pi/128$. Line with square: model coefficient; Line with delta: test coefficient.



FIG. 5: (Color online) Energy spectra (filter-scale $\Delta = 3\pi/64$) for decaying isotropic turbulence (a posteriori), at $t = 0$, $6\tau_0$ and $12\tau_0$, where τ_0 is the initial large eddy turnover time scale. Bold solid line: DNS; Dashed line: (a) CDSM1, (b) CDSM2, (c) SDSM, and (d) DSM.

CDSM2) can improve the LES results of isotropic turbulence, generally.

Then, let's see the case of mixed model. In Fig. 7, we present the energy spectra of three type mixed models (CDMM1, CDMM2, DMM) and DSM model at different filter scale, (a) $\Delta = 3\pi/64$ (b) $\Delta = 3\pi/128$. In (a), the spectrum of CDMM1 shows closer to the DNS spectrum, and CDSM1 and DSM show the similar trend, and are better than DMM in most wave-number. In (b) CDMM1 and CDMM2 predict the energy spectra better than other models particularly in high wave-number.



FIG. 6: (Color online) Energy spectra (filter-scale $\Delta = 3\pi/128$) for decaying isotropic turbulence (a posteriori), at $t = 0$, $6\tau_0$ and $12\tau_0$, where τ_0 is the initial large eddy turnover time scale. Bold solid line: DNS; Dashed line: (a) CDSM1, (b) CDSM2, (c) SDSM, and (d) DSM.



FIG. 7: (Color online) Energy spectra and for steady forced isotropic turbulence (a posteriori). Solid line: DNS; dashed line: CDMM1; DashDD line: CDMM2; line with diamond: DMM; line with delta: DSM. (a) filter scale $\Delta = 3\pi/64$ (b) filter scale $\Delta = 3\pi/128$.

In Fig. 8, we show the distribution of SGS dissipation with δ/η for the four LES models (CDMM1 ,CDMM2, DMM and DSM) and DNS, where δ is the consecutive filter scale and η is the Kolmogrov scale. From Fig. 8, we can see that the SGS energy dissipation of CDMM1 and CDMM2 are much closer to the DNS' than that of DMM and DSM in most of the range.

In Fig. 9, we display the probability density functions (PDF) of the four LES models at the filter scale $\Delta = 3\pi/64$, and the PDF of DNS is taken as comparison. As the limitation of DSM model, it can not predict the backscatters. Here, we can see that CDMM1, CDMM2 and DMM can capture the backscatters. Particulallywe also can be easy to see the PDFs of CDMM1 and



FIG. 8: (Color online) The SGS energy dissipation versus δ/η (a priori). Bold line: DNS; Line with square: CDMM1; Line with delta: CDMM2; Line with diamond: DMM; Line with circle: DSM.



FIG. 9: (Color online) Different probability density functions for the energy flux of DNS and other SGS models at the filter scale $\Delta = 3\pi/64$ (a posteriori) Bold line: DNS; Dashed line: CDMM1; Line with square: CDMM2; Line with delta: DMM; DashDD line: DSM.

CDMM2 fit the PDF of DNS very well. Through a priori test, we can get the results that model stress weight τ_{12}^{mod} of CDMM1, CDMM2 and DMM have a high correlation with the real stress weight τ_{12} over 85%, and the result of DSM has a low correlation with the real case not more than 30%.

We also display the distribution of high-order velocity increment with r/Δ in Fig. 10, where r is the separation distance and Δ is the filter scale $\Delta = 3\pi/64$. From the figure (a) (b) (c), we can see that CDMM1, CDMM2 and DSM are all close to DNS very well with no obvious difference in the range $r/\Delta \geqslant 3$. While in range of $r/\Delta \leqslant 3$, CDMM1, CDMM2 display better than DSM, but are still far from the result of DNS. In general, CDMM1, CDMM2 show better than DSM and DMM.



FIG. 10: (Color online) The distribution of high-order moments of longitudinal velocity increment with the separation distance r, where Δ is the filter scale. (a) S_4^L, (b) S_6^L, (c) S_8^L.

V. CONCLUSION

In this paper, we first put forward a general LES method to simulate the turbulent flows. Using the method, we can ignore the assumption of scale invariance, and we can also do LES out of the inertial range. The main idea of the method is construct a relation of SGS dissipation and viscous dissipation between two scale, and we use such a relationship as an additional constraint condition in the dynamic models to do large eddy simulation.

Through a priori and a posteriori tests, the single model CDSM1 and CDSM2 can predict energy spectra well in different range, they give the more proper model and test model coefficients. While the mixed model CDMM1 and CDMM2 also can predict energy spectra well in different range, and at the same time they predict the SGS energy dissipation more accurate, the PDF of energy flux much closer to the real case, and capturing the energy backscatter very well. Using the two models, the model stress has a higher correlation with the real SGS stress, and predicting the structure of the turbulence has a little progress.

Contrast with other constrained LES method, our method gives up the character of scale invariance and could also do LES well out of inertial range and even apply into simulating other type turbulence, such as turbulence in the boundary lay, compressible turbulence, etc.

---

[1] J. W. Deardoff, "The use of subgrid transport equation in a three dimensional model of atmospheric turbulence," ASME J. **26**(6), 669 (1973).

[2] J. Smagorinsky, "General circulation experiments with primitive equation," Monthly Weather Review, **91**, 99 (1963).

[3] P. Moin and J. Kim, "Numerical investigation of turbulent channel flow," J. Fluid Mech. **118**, 341 (1982).

[4] M. Germano, U. Piomelli, P. Moin, and W. Cabot, "A dynamic subgridscale eddy viscosity model," Phys. Fluids A **3**, 1760 (1991).

[5] D. K. Lilly, "A proposed modification of the Germano subgrid-scale closure method," Phys. Fluids A **4**, 633 (1992).

[6] C. meneveau and J. Katz, "Scale-invariance and turbulence models for large-eddy simulation," Annu. Rev. Fluid Mech. **32**, 1 (2000).

[7] S. B. Pope, *Turbulent Flows*, (Cambridge University Press, Cambridge, 2000).

[8] Y. Shi, Z. Xiao and S. Chen, "Constrained subgrid-scale stress model for large eddy simulation," Phys. Fluids **20**, 011701 (2008).

[9] S. C. C. Bailey, M. Hultmark, J. Schumacher, V. Yakhot, and A. J. Smits, "Measurement of local dissipation scales in turbulent pipe flow," Phys. Rev. Lett. **103**,014502 (2009).

[10] K. R. Sreenivasan and R. A. Antonia, "The phennomenology of small-scale turbulence," Ann. Rev. Fluid Mech. **29**, 435 (1997).

[11] L. Biferale, "A note on the fluctuation of dissipative scale in turbulence," Phys. Fluids **20**, 031703 (2008).

[12] M. Fatica, P. Orlandi, and R. Verzicco, "Direct and Large Eddy Simulations of Round Jets," Fluid Mechanics and Its Applications **26**, 49 (1994).

[13] C. Ronchi, M. Ypma, and V. M. Canuto, "On the application of the Germano identity to subgrid-scale modeling," Phys. Fluids A **12**, 2927 (1992).

[14] P. Voke, "Subgrid-scale modelling at low mesh Reynolds number," Theor. Comput. Fluid Dyn. **8**, 131 (1996).

[15] C. Meneveau, and T. S. Lund, "The dynamic Smogorinsky model and scale dependent coefficients in the viscous range of turbulence," Phys. Fluids **9**, 3932 (1997).

[16] C. Meneveau and T. S. Lund, "Dynamic model with scale-dependent coefficients in the viscous range," in Proceedings of the 1996 SummerProgram, Center for Turbulence Research, Stanford University, (1996).

[17] C. Meneveau, T. S. Lund, and W. Cabot, "A Lagrangian dynamic subgridscale model of turbulence,"

J. Fluid Mech. **319**, 353 (1996).

[18] F. Porte-Agel, C. Meneveau, and M. Parlange, "A scale-dependent dynamic model for large-eddy simulation: application to a neutral atmospheric boundary layer," J. Fluid Mech. **415**, 261 (2000).

[19] Elie B-Z, C. Meneveau, and M. Parlange, "A scale-dependent Lagrangian dynamic model for large eddy simulation of complex turbulent flows," Phys. Fluids **17**, 025105 (2005).

[20] W. Heisenberg, "On the theory of statistical and isotropic turbulence," Proc. Roy. Soc. London Ser. A, **195**, 402-406 (1948).

[21] Y. H. Pao, "Structure of turbulent velocity and scalar fields at large wavenumbers," Phys. Fluids **8**, 1063-1075 (1965)

[22] Z. S. She, "On the universal form of energy spectra in fully developed turbulence," Phys. Fluids A **5**, 1526-1528 (1993)

[23] S. Y. Chen, G. Doolen, J. R. Herring, R. H. Kraichnan, S. A. Orszag, and Z. S. She, "Far-dissipation range of turbulence," Phys. Rev. Lett. **70**, 3051-3054 (1993)

[24] L. S. G. Kovasnay, "spectrum of locally isotropic turbulence," J. Aero. Sci. **15**, 745 (1948).

[25] U. Frisch, P.L.Sulum, and M. Nelkin, "A simple dynamical model of intermittent fully developed turbulence," J. Fluid Mech. **87**, 719 (1978).

[26] H. Lu, C. J. Rutland, and L. M. Smith, "A priori ltests of one-equation LES modeling of rotating turbulence," Journal of Turbulence **37**, 1468 (2007).

[27] V. Borue and S. Orszag, "Local energy flux and subgrid-scale statistics in three-dimensional turbulence," J. Fluid Mech. **336**, 1 (1998).

[28] R. A. Clark, J. H. Ferziger, and W. C. Reynolds, "Evaluation of subgrid-scale models using an accurately simulated turbulent flow," J. fluid Mech. **91**, 1 (1979).

[29] S. Y. Chen, G. D. Doolen, R. H. Kraichnan, and Z. S. She, "on statistical correlations between velocity increments and locally averaged dissipation in homogeneous turbulence," Phys. Fluids A **5**, 458 (1993).